


Concanamycin A (high purity)



1mg

Rating: Not Rated Yet

Price

Sales price 7700 ?

Discount

[Ask a question about this product](#)

Manufacturer [BioViotica](#)

Description Product Details

Synonyms	Folimycin; Antibiotic TAN 1323B; Antibiotic X4357B
Product Type	Chemical
Properties	
Formula	C ₄₆ H ₇₅ NO ₁₄
MW	866.1
CAS	80890-47-7
RTECS	CB9732000
Source/Host Chemicals	Isolated from Streptomyces sp.
Purity Chemicals	?98% (HPLC)
Appearance	White to off-white solid.
Solubility	Soluble in methanol, DMSO or acetonitrile; insoluble in water.
Identity	Determined by 1H-NMR.
Declaration	Manufactured by BioViotica.
InChi Key	DJZCTUVALDDONK-MSPZYORZSA-N
Shipping and Handling	
Shipping	AMBIENT
Short Term Storage	+4°C
Long Term Storage	-20°C
Handling Advice	Protect from light when in solution.
Use/Stability	Stable for at least 1 year after receipt when stored at -20°C. After reconstitution protect from light at -20°C.

Antibiotic.

More potent and specific H⁺-ATPase inhibitor than bafilomycin A1 (Prod. No. BVT-0252).

Inhibits acidification of organelles such as lysosomes and the Golgi apparatus.

Inhibitor of autophagic degradation by rising lysosomal pH and thus inactivating the lysosomal acid hydrolases.

Uncategorized : Concanamycin A (high purity)

Blocks cell surface expression of viral glycoproteins without affecting their synthesis.
Cytotoxic in a number of cell lines in a cell viability assay.
Induces nitric oxide (NO) production.

Product References

Isolation and characterization of concanamycins A, B and C: H. Kinashi, et al.; *J. Antibiot.* 37, 1333 (1984)
Folimycin (concanamycin A), a specific inhibitor of V-ATPase, blocks intracellular translocation of the glycoprotein of vesicular stomatitis virus before arrival to the Golgi apparatus: M. Muroi, et al.; *Cell Struct. Funct.* 18, 139 (1993)
Folimycin (concanamycin A), an inhibitor of V-type H(+)-ATPase, blocks cell-surface expression of virus-envelope glycoproteins: M. Muroi, et al.; *BBRC* 193, 999 (1993)
Inhibitory effect of modified bafilomycins and concanamycins on P- and V-type adenosinetriphosphatases: S. Drose, et al.; *Biochemistry* 32, 3902 (1993)
Involvement of the vacuolar H(+)-ATPases in the secretory pathway of HepG2 cells: M. Yilla, et al.; *J. Biol. Chem.* 268, 19092 (1993)
Characterization of the ATPase activity of P-glycoprotein from multidrug-resistant Chinese hamster ovary cells: F.J. Sharom, et al.; *Biochem. J.* 308 (Pt2), 381 (1995)
Specific inhibitors of vacuolar type H(+)-ATPases induce apoptotic cell death: T. Nishihara, et al.; *BBRC* 212, 255 (1995)
Concanamycin A, the specific inhibitor of V-ATPases, binds to the V(o) subunit c: M. Huss, et al.; *J. Biol. Chem.* 277, 40544 (2002)
Nitric oxide production by the vacuolar-type (H+)-ATPase inhibitors bafilomycin A1 and concanamycin A and its possible role in apoptosis in RAW 264.7 cells: J. Hong, et al.; *J. Pharmacol. Exp. Ther.* 319, 672 (2006)
Degradation of oxidized proteins by autophagy during oxidative stress in Arabidopsis: Y. Xiong, et al.; *Plant Physiol.* 143, 291 (2007)
Inhibitors of the V0 subunit of the vacuolar H+-ATPase prevent segregation of lysosomal- and secretory-pathway proteins: J.A. Sobota, et al.; *J. Cell Sci.* 122, 3542 (2009)
Inhibitors of V-ATPase: old and new players: M. Huss, et al.; *J. Exp. Biol.* 121, 341 (2009)
Increased production of reactive oxygen species by the vacuolar-type (H+)-ATPase inhibitors bafilomycin A1 and concanamycin A in RAW 264 cells: A. Yokomakura, et al.; *J. Toxicol. Sci.* 37, 1045 (2012)
Inhibition of vacuolar ATPase attenuates the TRAIL-induced activation of caspase-8 and modulates the trafficking of TRAIL receptosomes: V. Horova, et al.; *FEBS J.* 280, 3436 (2013)
Estimating the rotation rate in the vacuolar proton-ATPase in native yeast vacuolar membranes: Z. Ferencz, et al.; *Eur. Biophys. J.* 42, 147 (2013)
Inhibitors of vacuolar ATPase proton pumps inhibit human prostate cancer cell invasion and prostate-specific antigen expression and secretion: V. Michel, et al.; *Int. J. Cancer* 132, E1 (2013)
Loss of vacuolar H+-ATPase activity in organelles signals ubiquitination and endocytosis of the yeast plasma membrane proton pump Pma1p: A. M. Smardon & P. M. Kane; *J. Biol. Chem.* 289, 32316 (2014)
Combined effects of EGFR tyrosine kinase inhibitors and vATPase inhibitors in NSCLC cells: H. Jin, et al.; *Toxicol. Appl. Pharm.* 278, 17 (2015)
Appropriate vacuolar acidification in *Saccharomyces cerevisiae* is associated with efficient high sugar fermentation: T.D. Nguyen, et al.; *Food Microbiology* 70, 262 (2018)

Reviews

There are yet no reviews for this product.